

simass.txt

This is a file with simulation assessment, summarising performance criteria over model domain. The file is located in the [resultdir](#) folder. The file contains values of most [performance criteria](#) of the selected variables in the objective function. Note: If several RA criteria have been selected, only the last of them will be printed to file. Not calculated criterion are indicated by -9999. All information from the simass-files can also be found in the [hyss_yymmddHHMM.log](#).

When ensemble or sequence simulations are made, the results from simulations ($l=1 \dots n$ or $l=\text{sequence number}>0$) are written to files named simassX_00l.txt, where n is defined by num_ens in [optpar.txt](#).

For the calculation of criterion for lake water stage, the combination of variables wcom and wstr are exchanged for the internal variables clwc and clws by the program. These variables are the water stages cleaned from w0ref reference level ($clwc=wcom-w0ref$, $clws=wstr-w0ref$). This makes the criterion calculation more accurate, but note that relative criteria, e.g. relative bias, are relative to the smaller cleaned water stage level.

The following performance criteria may be calculated: Code is corresponding code for [info.txt](#). Equations can be found here.

Criterion	Code	Description
Regional NSE	RR2	regional Nash-Sutcliffe efficiency (all data combined in one long time-series)
Regional RA	RRA	regional Nash-Sutcliffe efficiency like criteria where the square is exchanged with a coefficient value
Regional RE	RRE	regional relative bias (all data combined in one long time-series)
Regional MAE	-	regional absolute error (all data combined in one long time-series)
Average NSE	MR2	average of Nash-Sutcliffe efficiencies for all subbasins with observations
Average RA	MRA	average value of subbasin values of Nash-Sutcliffe like criteria where the square is exchanged with a coefficient value
Average RE	MRE	average of the relative bias for all subbasins (Note: fraction, not %)
Average RSDE	MRS	error in standard deviation, average of all subbasins with observations
Average CC	MCC	Pearson correlation coefficient, average of all subbasins with observations
Average ARE	MAR	mean absolute of relative errors for all subbasins (Note: fraction, not %)
Spatial NSE	SR2	spatial R2 calculated using annual means for all subbasins (requires at least 5 years and 5 subbasins with data)
Spatial RA	RRA	Spatial Nash-Sutcliffe like criteria where the square in the Nash-Sutcliffe formula is exchanged for a coefficient value
Spatial RE	-	spatial relative error calculated using annual means for all subbasins (requires at least 5 years and 5 subbasins with data)
Kendalls Tau	TAU	average of Kendall's Tau value for all subbasins
Median NSE	MD2	median of Nash-Sutcliffe efficiency for all subbasins with observations
Median RA	MDA	median of all subbasins RA (Nash-Sutcliffe like criteria where the square is exchanged with a coefficient value)
Median KGE	MKG	median of all subbasins Kling-Gupta efficiency
Median NRMSE	MNR	median of all subbasins normalised RMSE

